



All orders must be reviewed, completed and signed by the prescriber before they will be implemented.
To delete an order that does not have a 'check box' preceding it the order must be stroked out and initialed. For orders where check boxes are provided, the prescriber indicates with a check mark to confirm the order; if left blank, the order will not be activated.

Page No. _____	Allergies: <input type="checkbox"/> NO <input type="checkbox"/> YES (specify reaction): _____
Pain Management PRN Orders	
Date: _____	Time: _____
MD Instructions: PRN opioid medication and rescue dose opioids may NOT be given if the patient is already on PCA or epidural analgesia unless approved by the Pain Service.	
Assessment, Monitoring & Management:	
<input checked="" type="checkbox"/> Assess and monitor patient and titrate opioid as per policies "Opioid Titration Policy" and "Policy for the Assessment, management, and monitoring of patients prescribed opioids and/or local anesthetics"	
Acetaminophen:	
<input type="checkbox"/> Acetaminophen 1000 mg PO q6h x _____ days (maximum dose from all sources must not exceed 4 g per 24 h) or <input type="checkbox"/> Acetaminophen 650 mg PO q6h x _____ days (maximum dose from all sources must not exceed 4 g per 24 h) <input type="checkbox"/> Acetaminophen 650 mg PR q6h x _____ days (while awake) if unable to take PO meds (maximum dose from all sources must not exceed 4 g per 24 h) <input type="checkbox"/> Acetaminophen 650 mg PO q6h PRN x _____ days (maximum dose from all sources must not exceed 4 g per 24 h)	
Nonsteroidal anti-inflammatory drugs (Choose 1):	
MD Instructions: Not applicable if patient already taking a non-steroidal anti-inflammatory drug Indomethacin not recommended for postpartum patients Misoprostol contraindicated for obstetrical patients Caution in patients with history of or risk of GI bleed, renal insufficiency, renal transplant; do not use if creatinine clearance less than 30mL/min	
<input type="checkbox"/> Indomethacin 50 mg PO/PR q8h for _____ days or <input type="checkbox"/> Naproxen 500 mg PO/PR q12h for _____ days or <input type="checkbox"/> Ibuprofen 400 mg PO q6h for _____ days or <input type="checkbox"/> Ibuprofen 400 mg PO q8h for _____ days (for patients over age 75) or <input type="checkbox"/> Meloxicam 7.5 mg PO q24h for _____ days or <input type="checkbox"/> Meloxicam 15 mg PO q24h for _____ days or <input type="checkbox"/> Other (Specify): _____ for _____ days	
AND CONSIDER:	
<input type="checkbox"/> Misoprostol 200 mcg PO QID for _____ days while on non-steroidal anti-inflammatory drug or <input type="checkbox"/> Pantoprazole 40 mg PO once daily for _____ days while on non-steroidal anti-inflammatory drug or <input type="checkbox"/> Other (Specify): _____	
PRN Opioid medications for pain (Choose 1):	
MD Instructions: When prescribing IV analgesia for moderate pain - consider lower opioid doses for elderly patients, patients with low body weight and severe co-morbidities – in moderate to severe pain, starting IV dose in these patients is <i>usually</i> within range of morphine 0.05 – 0.1 mg/kg or hydromorphone 0.0075 – 0.015 mg/kg per dose. MD to reassess opioid medication orders if: <ul style="list-style-type: none"> • Patient has taken more than 6 PRN doses (maximum doses) in past 24hrs • Patient has persistent pain (moderate or severe or pain scale greater than or equal to 4/10) after two doses of PRN Opioid 	

Print Name: _____ Signature: _____ **MD** _____

All orders are to be signed including Qualifying Initials



All orders must be reviewed, completed and signed by the prescriber before they will be implemented.
To delete an order that does not have a 'check box' preceding it the order must be stroked out and initialed. For orders where check boxes are provided, the prescriber indicates with a check mark to confirm the order; if left blank, the order will not be activated.

Page No. _____	Allergies: <input type="checkbox"/> NO <input type="checkbox"/> YES (specify reaction):
Pain Management PRN Orders	
medications	Transcribed
Date: _____ Time: _____	
<ul style="list-style-type: none"> • Patient has intolerable side effects • Patient has increased sedation (rousable but frequently drowsy, and drifts off to sleep during conversation, or somnolent or has minimal or no response to stimulus) • Signs of respiratory depression (respiratory rate less than 10/minute, periods of apnea / irregular breathing or airway obstruction) 	
<p>For Mild pain(Choose 1):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Morphine 5-10 mg PO q4h PRN or Morphine 3 - 5 mg subcut or IV q4h PRN (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> HYDROMorphone 2 - 4 mg PO q4h PRN or HYDROMorphone 0.5 – 1 mg subcut or IV q4h PRN (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> Tylenol #3 (use Acetaminophen 300 mg with codeine 30 mg) 1-2 Tabs PO q4h PRN (maximum dose of acetaminophen from all sources must not exceed 4 g per 24 h (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> Other (Specify): _____ (Assess and monitor patient and titrate opioid as per policy) <p>For Moderate to Severe pain (Choose 1):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Acetaminophen 325 mg with oxycodone 5 mg (Percocet) 1-2 Tabs PO q4h PRN (maximum dose of acetaminophen from all sources must not exceed 4 g per 24 h) (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> Oxycodone IR (immediate release) 10 – 15 mg PO q4h PRN (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> Morphine 10-15 mg PO q4h PRN or Morphine 5 - 7.5 mg subcut or IV q4h PRN (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> Morphine 15-20 mg PO q4h PRN or Morphine 7.5 -10 mg subcut or IV q4h PRN (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> HYDROMorphone 4 - 6 mg PO q4h PRN or HYDROMorphone 1 - 1.5 mg subcut or IV q4h PRN (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> HYDROMorphone 6 - 8 mg PO q4h PRN or HYDROMorphone 1.5 – 2 mg subcut or IV q4h PRN (Assess and monitor patient and titrate opioid as per policy) <input type="checkbox"/> Other (Specify): _____ (Assess and monitor patient and titrate opioid as per policy) 	
AND CONSIDER:	
<ul style="list-style-type: none"> <input type="checkbox"/> DiphenhydrAMINE (Benadryl) 25 mg IV/PO q4h PRN for pruritus <input type="checkbox"/> Ondansetron 4mg IV/PO q6h PRN for nausea <input type="checkbox"/> DimenhydrINATE (Gravol) 25 mg IV or 50 mg PO q4h PRN for nausea <input type="checkbox"/> Metoclopramide 10 mg IV q4h PRN for nausea (not to exceed 40 mg/day) <input type="checkbox"/> Docusate sodium 100 mg PO BID <input type="checkbox"/> Sennosides 17.2 mg po qhs PRN for constipation <input type="checkbox"/> Lactulose 15-30 mL po bid PRN for constipation 	

Print Name: _____ Signature: _____ **MD**

All orders are to be signed including Qualifying Initials